IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Mplicad (S): Sames A. Thomson

erial**/Ng/:** 08/376,327

anuary 20, 1995

PRIMATE EMBRYONIC STEM CELLS

Art Unit:

Examiner:

Atty Docket:

960296.92905

Date:

October 7, 1995

HEC WED

NOV G 2 1995

GROUT 1800

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

Transmitted herewith is an amendment in the above-identified patent application.

The fee for that amendment has been calculated as shown below:

CLAIMS AS AMENDED

	Claims After Amendment		Highest Number Paid For Previously	Numbe	er Extra	Rate	Additional Fee
Total Claims		Minus		0	X	\$11.00	= \$.00
Independent Claims		Minus		0	Х	\$39.00	= \$.00
First presentation of a Multiple Dependent Claim				\$125.00	= \$.00		
						Total Fee	\$.00

- [X] No additional fee is required.
- [] A check for \$.00 to cover the filing fee and the cost of recording the assignment is enclosed.
- [X] The Commissioner is hereby authorized to charge any additional fees which may be required or credit any overpayment to Account No. 17-0055. An extra copy of this sheet is enclosed.

Respectfully submitted,

Nicholas J. Seay

Registration No. 27,386

Quarles and Brady P.O. Box 2113 Madison, WI 53701-2113 (608) 251-5000



435/373

United States Patent [19]

US003. SOA

[11] Patent Number:

5,843,780

[45] Date of Patent:

Dec. 1, 1998

[54] PRIMATE EMBRYONIC STEM CELLS

[75] Inventor: James A. Thomson, Madison, Wis.

[73] Aszignee: Wisconsin Alumni Research

Foundation, Madison, Wis.

[21] Appl. No.: 591,246

Thomson

[22] Filed: Jan. 18, 1996

Related U.S. Application Data

[63]	Continuation-in-part of Ser. No. 376,327, Jan. 20, 1995.
[51]	Int. CL ⁶ C12N 5/06
	U.S. Cl 435/363; 435/366; 435/373
[58]	Field of Search

[56] References Cited

U.S. PATENT DOCUMENTS

5,449,620	9/1995	Khillan .	
5,453,357	9/1995	Hogan	435/7.21
5,591,625	1/1997	Gerson et al	

FOREIGN PATENT DOCUMENTS

WO 94/03585 2/1994 WIPO .

OTHER PUBLICATIONS

Bongso, et al., "Isolation and culture of inner cell mass cells from human blastocysts", Human Reproduction, 9:2110-2117, 1994.

Brown, et al., "Criteria that optimize the potential of murine embryonic stem cells for in vitro and in vivo developmental studies", In Vitro Cell. Dev. Biol. 284:773–778, Dec. 1992.

Damjanov, et al., "Retinoic acid-induced differentiation of the developmentally pluripotent human germ cell tumordervied cell line, NCCIT", Laboratory Investigation, 68:220-232, 1993.

Nation/World, "Embryonic monkey cells isolated". -Nov. 4, 1994.

Bongso, A., et al., "The Growth of Inner Cell Mass Cells from Human Blastocysts," *Theriogenology*, 41:167 (1994).

Thomson, James A., et al., "Pluripotent Cell Lines Derived from Common Marmoeet (Callithrix jacchus) Blastocysts," Biology of Reproduction, 55:254-259 (1996).

Primary Examiner—Michael P. Woodward Assistant Examiner—Brenda G. Brumback Attorney, Agent, or Firm—Quaries & Brady

[57] ABSTRACT

A purified preparation of primate embryonic stem cells is disclosed. This preparation is characterized by the following cell surface markers: SSEA-1 (-); SSEA-3 (+); SSEA-4 (+); TRA-1-60 (+); TRA-1-81 (+); and alkaline phosphatase (+). In a particularly advantageous embodiment, the cells of the preparation have normal karyotypes and continue to proliferate in an undifferentiated state after continuous culture for eleven months. The embryonic stem cell lines also retain the ability, throughout the culture, to form trophoblast and to differentiate into all tissues derived from all three embryonic germ layers (endoderm, mesoderm and ectoderm). A method for isolating a primate embryonic stem cell line is also disclosed.

11 Claims, 8 Drawing Shoots





Thomson

[11] Patent Number:

5,843,780

[45] Date of Patent:

Dec. 1, 1998

[54]	PRIMATE	EMBRYONIC	STEM	CELLS
------	---------	------------------	-------------	-------

[75] Inventor: James A. Thomson, Madison, Wis.

[73] Assignee: Wisconsin Alumni Research Foundation, Madison, Wis.

[21] Appl. No.: 591,246

[22] Filed: Jan. 18, 1996

Related U.S. Application Data

[63]	Continuation-in-part of	Ser. No. 376,327, Jan. 20, 1995.
[51]	Int. Cl.6	C12N 5/06
[52]	U.S. Cl	 435/363 ; 435/366; 435/373
[58]	Field of Search	435/363, 366,
- 1		435/373

[56]

References Cited

U.S. PATENT DOCUMENTS

5,449,620	9/1995	Khillan .	
5,453,357	9/1995	Hogan	435/7.21
5,591,625	1/1997	Gerson et al	

FOREIGN PATENT DOCUMENTS

WO 94/03585 2/1994 WIPO .

OTHER PUBLICATIONS

Bongso, et al., "Isolation and culture of inner cell mass cells from human blastocysts", Human Reproduction, 9:2110-2117, 1994.

Brown, et al., "Criteria that optimize the potential of murine embryonic stem cells for in vitro and in vivo developmental studies", In Vitro Cell. Dev. Biol. 284:773-778, Dec. 1992.

Damjanov, et al., "Retinoic acid-induced differentiation of the developmentally pluripotent human germ cell tumordervied cell line, NCCIT", Laboratory Investigation, 68:220-232, 1993.

Nation/World, "Embryonic monkey cells isolated". -Nov. 4, 1994

Bongso, A., et al., "The Growth of Inner Cell Mass Cells from Human Blastocysts," *Theriogenology*, 41:167 (1994).

Thomson, James A., et al., "Phuripotent Cell Lines Derived from Common Marmoset (Callithrix jacchus) Blastocysts," Biology of Reproduction, 55:254-259 (1996).

Primary Examiner—Michael P. Woodward Assistant Examiner—Brenda G. Brumback Attorney, Agent, or Firm—Quarles & Brady

[57]

ABSTRACT

A purified preparation of primate embryonic stem cells is disclosed. This preparation is characterized by the following cell surface markers: SSEA-1 (-); SSEA-3 (+); SSEA-4 (+); TRA-1-60 (+); TRA-1-81 (+); and alkaline phosphatase (+). In a particularly advantageous embodiment, the cells of the preparation have normal karyotypes and continue to proliferate in an undifferentiated state after continuous culture for eleven months. The embryonic stem cell lines also retain the ability, throughout the culture, to form trophoblast and to differentiate into all tissues derived from all three embryonic germ layers (endoderm, mesoderm and ectoderm). A method for isolating a primate embryonic stem cell line is also disclosed.

11 Claims, 8 Drawing Sheets

DKT-5/1169-1016 LIZGOIN 912-863-2086 Forg 2865





Thomson

[11] Patent Number:

5,843,780

[45] Date of Patent:

Dec. 1, 1998

[54] PRIMATE EMBRYONIC STEM CELLS

[75] Inventor: James A. Thomson, Madison, Wis.

[73] Assignee: Wisconsin Alumni Research

Foundation, Madison, Wis.

[21] Appl. No.: 591,246

[22] Filed: Jan. 18, 1996

Related U.S. Application Data

[63] Continuation-in-part of Ser. No. 376,327, Jan. 20, 1995.

435/373

[56] References Cited

5,591,625

U.S. PATENT DOCUMENTS

FOREIGN PATENT DOCUMENTS

1/1997 Gerson et al. .

WO 94/03585 2/1994 WIPO.

OTHER PUBLICATIONS

Bongso, et al., "Isolation and culture of inner cell mass cells from human blastocysts", Human Reproduction, 9:2110-2117, 1994.

Brown, et al., "Criteria that optimize the potential of murine embryonic stem cells for in vitro and in vivo developmental studies", In Vitro Cell. Dev. Biol. 284:773-778, Dec. 1992.

Damjanov, et al., "Retinoic acid-induced differentiation of the developmentally pluripotent human germ cell tumordervied cell line, NCCIT", Laboratory Investigation, 68:220-232, 1993.

Nation/World, "Embryonic monkey cells isolated". -Nov. 4, 1994.

Bongso, A., et al., "The Growth of Inner Cell Mass Cells from Human Blastocysts," *Theriogenology*, 41:167 (1994).

Thomson, James A., et al., "Pluripotent Cell Lines Derived from Common Marmoset (Callithrix jacchus) Blastocysts," Biology of Reproduction, 55:254-259 (1996).

Primary Examiner—Michael P. Woodward Assistant Examiner—Brenda G. Brumback Attorney, Agent, or Firm—Quarles & Brady

[57] ABSTRACT

A purified preparation of primate embryonic stem cells is disclosed. This preparation is characterized by the following cell surface markers: SSEA-1 (-); SSEA-3 (+); SSEA-4 (+); TRA-1-60 (+); TRA-1-81 (+); and alkaline phosphatase (+). In a particularly advantageous embodiment, the cells of the preparation have normal karyotypes and continue to proliferate in an undifferentiated state after continuous culture for eleven months. The embryonic stem cell lines also retain the ability, throughout the culture, to form trophoblast and to differentiate into all tissues derived from all three embryonic germ layers (endoderm, mesoderm and ectoderm). A method for isolating a primate embryonic stem cell line is also disclosed.

11 Claims, 8 Drawing Sheets





Thomson

[11] Patent Number:

5,843,780

[45] Date of Patent:

Dec. 1, 1998

[54]	PRIMATE EMBRYONIC STEM CELLS	Damja
		the de

[75] Inventor: James A. Thomson, Madison, Wis.

[73] Assignee: Wisconsin Alumni Research Foundation, Madison, Wis.

[21] Appl. No.: 591,246

[22] Filed: Jan. 18, 1996

Related U.S. Application Data

[63]	Continuation-in-	part of Ser. No	o. 376,327, Jan.	20, 1995.
------	------------------	-----------------	------------------	-----------

[51] Int. Cl.6 C12N 5/06

[52] U.S. Cl. 435/363; 435/366; 435/373

[56]

References Cited

U.S. PATENT DOCUMENTS

5,449,620	9/1995	Khillan .	
5,453,357	9/1995	Hogan	435/7.21
5,591,625	1/1997	Gerson et al	

FOREIGN PATENT DOCUMENTS

WO 94/03585 2/1994 WIPO .

OTHER PUBLICATIONS

Bongso, et al., "Isolation and culture of inner cell mass cells from human blastocysts", Human Reproduction, 9:2110-2117, 1994.

Brown, et al., "Criteria that optimize the potential of murine embryonic stem cells for in vitro and in vivo developmental studies", In Vitro Cell. Dev. Biol. 284:773-778, Dec. 1992.

Damjanov, et al., "Retinoic acid-induced differentiation of the developmentally pluripotent human germ cell tumordervied cell line, NCCIT", Laboratory Investigation, 68:220-232, 1993.

Nation/World, "Embryonic monkey cells isolated". -Nov. 4, 1994.

Bongso, A., et al., "The Growth of Inner Cell Mass Cells from Human Blastocysts," *Theriogenology*, 41:167 (1994).

Thomson, James A., et al., "Pluripotent Cell Lines Derived from Common Marmoset (Callithrix jacchus) Blastocysts," Biology of Reproduction, 55:254-259 (1996).

Primary Examiner—Michael P. Woodward Assistant Examiner—Brenda G. Brumback Attorney, Agent, or Firm—Quarles & Brady

[57]

ABSTRACT

A purified preparation of primate embryonic stem cells is disclosed. This preparation is characterized by the following cell surface markers: SSEA-1 (-); SSEA-3 (+); SSEA-4 (+); TRA-1-60 (+); TRA-1-81 (+); and alkaline phosphatase (+). In a particularly advantageous embodiment, the cells of the preparation have normal karyotypes and continue to proliferate in an undifferentiated state after continuous culture for eleven months. The embryonic stem cell lines also retain the ability, throughout the culture, to form trophoblast and to differentiate into all tissues derived from all three embryonic germ layers (endoderm, mesoderm and ectoderm). A method for isolating a primate embryonic stem cell line is also disclosed.

11 Claims, 8 Drawing Sheets





Thomson

[11] Patent Number:

5,843,780

[45] Date of Patent:

Dec. 1, 1998

[54] PRIMATE EMBRYONIC STEM CELLS

[75] Inventor: James A. Thomson, Madison, Wis.

[73] Assignee: Wisconsin Alumni Research

Foundation, Madison, Wis.

[21] Appl. No.: 591,246

[22] Filed: Jan. 18, 1996

Damjanov. et al.. "Retinoic acid-induced differentiation of the developmentally pluripotent human germ cell tumor-dervied cell line, NCCIT". Laboratory Investigation. 68:220-232. 1993.

Nation/World. "Embryonic monkey cells isolated". -Nov. 4, 1994.

Bongso, A., et al., "The Growth of Inner Cell Mass Cells from Human Blastocysts." *Theriogenology*, 41:167 (1994).

Thomson, James A., et al., "Pluripotent Cell Lines Derived from Common Marmoset (*Callithrix jacchus*) Blastocysts," *Biology of Reproduction*, 55:254-259 (1996).

ABSTRACT

Primary Examiner—Michael P. Woodward Assistant Examiner—Brenda G. Brumback Attorney, Agent, or Firm—Quarles & Brady

[57]

435/373

A purified preparation of primate embryonic stem cells is disclosed. This preparation is characterized by the following cell surface markers: SSEA-1 (-); SSEA-3 (+); SSEA-4 (+); TRA-1-60 (+); TRA-1-81 (+); and alkaline phosphatase (+). In a particularly advantageous embodiment, the cells of the preparation have normal karyotypes and continue to proliferate in an undifferentiated state after continuous culture for eleven months. The embryonic stem cell lines also retain the ability, throughout the culture, to form trophoblast and to differentiate into all tissues derived from all three embryonic germ layers (endoderm, mesoderm and ectoderm). A method for isolating a primate embryonic stem cell line is also disclosed.

11 Claims, 8 Drawing Sheets

Related U.S. Application Data

	Continuation-in-part of Ser. No	
[51]	Int. Cl.6	C12N 5/06
[52]	U.S. Cl 43	5/363 ; 435/366; 435/373
[58]	Field of Search	435/363, 366.

[56]

References Cited

U.S. PATENT DOCUMENTS

5,449,620	9/1995	Khillan .	
5,453,357	9/1995	Hogan	435/7.21
5,591,625	1/1997	Gerson et al	

FOREIGN PATENT DOCUMENTS

WO 94/03585 2/1994 WIPO .

OTHER PUBLICATIONS

Bongso, et al., "Isolation and culture of inner cell mass cells from human blastocysts". Human Reproduction, 9:2110-2117, 1994.

Brown. et al., "Criteria that optimize the potential of murine embryonic stem cells for in vitro and in vivo developmental studies". In Vitro Cell. Dev. Biol. 284:773–778, Dec. 1992.